

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>A61K 31/195, 31/47, 33/30, 31/325, 31/42, 31/13, 31/44</b>		<b>A2</b>	(11) International Publication Number: <b>WO 97/09976</b>
			(43) International Publication Date: 20 March 1997 (20.03.97)
(21) International Application Number: <b>PCT/IB96/00981</b>		<b>(81) Designated States:</b> AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 23 August 1996 (23.08.96)			
(30) Priority Data: 60/003,134 1 September 1995 (01.09.95) US 60/007,356 20 November 1995 (20.11.95) US			
(71) Applicant: WASHINGTON UNIVERSITY [US/US]; One Brookings Drive, Campus Box 8013, St. Louis, MO 63130 (US).			
(72) Inventors: CHOI, Dennis, Wonkyu; 7400 Forsyth Boulevard, St. Louis, MO 63105 (US). KOH, Jae-young; 8148 Flintstone Trail, St. Louis, MO 63123 (US).			
(74) Agent: POKRAS, Bruce, A.; 340 Kingsland Street, Nutley, NJ 07110-1199 (US).		<b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title: METHOD OF REDUCING NEUROTOXIC INJURY WITH ZINC CHELATORS</b>			
<b>(57) Abstract</b>  The invention relates to the use of pharmaceutically acceptable zinc chelating compounds for the manufacture of medicaments for the treatment of neurotoxic injury.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

Method of reducing neurotoxic injury with zinc chelators

The invention relates to the use of zinc chelating compounds in the treatment of neurotoxic injury.

It has been discovered that zinc chelating compounds protect  
5 hippocampal neurons from the neurotoxic effects of  $Zn^{2+}$  released during neurotoxic events, such as ischemia as a result of stroke or cardiac arrest. Thus, the present invention comprises a method of treating neurotoxic injury in a patient suffering said injury, comprising administering to said patient a zinc chelating compound in an amount sufficient to treat said  
10 neurotoxic injury.

The administration of the composition comprising the zinc chelating compound is carried out by injection or infusion of the composition into the patients cerebrospinal fluid.

Chelatable  $Zn^{2+}$  is present in large quantities in presynaptic vesicles of  
15 central excitatory neurons (Danscher et al., 1985; Frederickson et al., 1983), and released with synaptic activity or membrane depolarization (Assaf and Chung, 1984; Howell et al., 1984; Charton et al., 1985). Although the precise role of released synaptic zinc is not known, it blocks NMDA receptor-mediated current (Westbrook and Mayer, 1987; Peters et al., 1987; Christine  
20 and Choi, 1990) and GABA receptor-mediated current (Westbrook and Mayer, 1987), as well as voltage-dependent calcium channels (Winegar and Lansman, 1990). In addition, exposure to excessive extracellular  $Zn^{2+}$  is neurotoxic to cortical neurons (Choi et al., 1988), possibly mediated by  $Zn^{2+}$  influx in part mediated by ionotropic glutamate receptors and voltage-gated  
25 calcium channels (Weiss et al., 1993; Koh and Choi, 1994). Recently, it has been shown that zinc translocates into degenerating hippocampal hilar neurons after transient ischemia (Tonder et al., 1990) or kainate-induced seizures (Frederickson et al., 1989).

Treating neurotoxic injury within the meaning of the present invention  
30 means reducing the extent of damage to central neurons surrounding a central neuron which has released  $Zn^{2+}$  due to its having been damaged by a

neurotoxic event. Neurotoxic events include acute neurological insults such as hypoxia/ischemia, such as occurs during stroke, cardiac arrest, hypoglycemia, epilepsy or trauma. Neurotoxic events may also be chronic neuronal damage caused by neurodegenerative disorders such as

5 Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and the neurodegenerative effects of AIDS. Thus, the present invention also comprises a method of treating diseases, such as those described above, in which said neurotoxic injury occurs.

The zinc chelating compounds useful in accordance with the invention

10 are not critical. Any conventional compound which is capable of chelating  $Zn^{2+}$  and which is pharmaceutically acceptable for injection into cerebral spinal fluid may be used in accordance with the invention.

While ethylenediaminetetraacetic acid (EDTA) is traditionally used as an organic chelating agent for  $Ca^{2+}$ , it has affinity for other divalent metal

15 ions, and specifically binds  $Zn^{2+}$  with much higher affinity than  $Ca^{2+}$  (log stability constant at pH 7, 13.1 for  $Zn^{2+}$  versus 7.3 for  $Ca^{2+}$ ). An equimolar combination for  $Ca^{2+}$  and EDTA will act as a chelating agent for  $Zn^{2+}$  without chelating local  $Ca^{2+}$  (Dansher et al., 1975). Thus, the preferred zinc chelating compound for use in accordance with the invention is disodium-

20 calcium EDTA ("CaEDTA"), especially the form of CaEDTA known as edetate calcium disodium injection, USP (e.g., Calcium Disodium Versenate, 3M Pharmaceuticals, St. Paul, Minnesota).

Another preferred zinc chelating compound useful in accordance with the invention is 3-mercapto-D-valine (penicillamine), which is a chelating

25 agent usually applied to the treatment of Wilson's disease where it removes excess copper.

Other zinc chelating compounds useful in accordance with the invention are:

bis(diethylthiocarbamoyl) disulfide (Disulfiram or Antabuse),

30 (ethylenedioxy)diethylenedinitrilotetraacetic acid (EGTA),  
N,N,N',N'-tetrakis(2-pyridylmethyl)-ethylenediamine (TPEN),  
N-(6-methoxy-8-quinolyl)-p-toluenesulfonamide (TSQ),  
8-hydroxy quinoline (Oxyne),  
8-hydroxy quinoline-5-sulphonic acid (Sulphoxine),

35 diethyl dithiocarbamate (DEDTC),

- 1,10-phenanthroline,  
dipicolinate,  
N-acetyl cystein,  
diphenylthiocarbazone (Dithizone),  
5 1-[2-(5-carboxyoxazol-2-yl)-6-aminobenzofuran-5-oxy]-2-(2'-amino-5'-methylphenoxy)ethane-N,N,N',N'-tetraacetic acid (Fura-2), and  
1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (BAPTA).

The zinc chelating compound used in accordance with the present invention may be administered to the cerebrospinal fluid by any conventional  
10 means. The preferred method of administration is by injection or chronic pump infusion into the lateral ventricles of the patient's brain or into the lumbar sac of the patient.

The zinc chelating compound is preferably administered as a composition containing the zinc chelating compound and a  
15 pharmaceutically acceptable carrier compatible with the compound. In preparing such a composition, any conventional pharmaceutically acceptable carrier may be utilized. Typical carriers for administration by injection would be sterile aqueous/buffered solutions, preferably water for injection or unbuffered or buffered physiological saline. The zinc chelating  
20 compound is preferably present in the carrier at a concentration of 1 mM - 300 mM, especially from 50 mM - 200 mM.

In carrying out the method of the invention, the zinc chelating compound is administered to adults daily in an amount from about 0.1 mg/kg to about 100 mg/kg daily, in single or divided doses, or continuously  
25 through chronic pump infusion. The preferred dosage will vary depending upon the indication for which the method of the invention is being used to treat. For treatment of patients who have suffered an acute neurotoxic insult, as a result of, e.g., stroke or cardiac arrest, the administration of a dosage from about 5 mg/kg to about 50 mg/kg daily, carried out for from 1 to 7  
30 days, is preferred. Administration may be carried out by injection or by infusion utilizing a chronic infusion pump. For treatment of chronic neurotoxic conditions in patients, e.g., Alzheimer's disease, a dosage of from about 1 mg/kg to about 10 mg/kg daily is preferred, carried out for months to years, preferably utilizing a chronic infusion pump.

35 Thus, the present invention also comprises a method of treating neurotoxic injury in a patient suffering said injury by administering to said

patient a composition comprising a zinc chelating compound and carrier, wherein both the compound and the carrier are pharmaceutically acceptable for injection. The preferred zinc chelating compounds are CaEDTA and 3-mercapto-D-valine, with CaEDTA being especially preferred. The zinc  
5 chelating compound is preferably administered in an amount from about 0.1 mg/kg to about 100 mg/kg daily, especially in an amount from about 5 mg/kg to about 50 mg/kg daily for the treatment of acute neurotoxic conditions and in an amount from about 1 mg/kg to about 10 mg/kg daily for chronic  
10 neurotoxic conditions. The preferred method of administration is the injection or infusion of the pharmaceutical composition comprising the zinc chelating compound and the carrier into the cerebrospinal fluid (e.g., the lateral ventricles or lumbar sac) of the patient. The preferred duration of treatment is from 1 to 7 days for acute neurotoxic conditions, and for months to years for chronic neurotoxic conditions.

15 The present invention preferably comprises a method of treating stroke in a patient suffering said stroke by administering to said patient a composition comprising a zinc chelating compound and a carrier, wherein both the compound and the carrier are pharmaceutically acceptable for injection. The preferred zinc chelating compounds are CaEDTA and 3-  
20 mercapto-D-valine, with CaEDTA being especially preferred. The zinc chelating compound is preferably administered in an amount from about 0.1 mg/kg to about 100 mg/kg daily, especially in an amount from about 5 mg/kg to about 50 mg/kg daily for the treatment of stroke. The preferred method of administration is the injection or infusion, especially injection, of the  
25 pharmaceutical composition comprising the zinc chelating compound and the carrier into the cerebrospinal fluid (e.g., the lateral ventricles or lumbar sac) of the patient. The preferred duration of treatment is from 1 to 7 days.

Experiments were conducted which determined that a zinc chelating compound, CaEDTA, exhibited neuroprotective effects in cell culture and  
30 animal models of brain hypoxic-ischemic injury, such as might occur consequent to cardiac arrest or stroke. Several observations were made:

1) In mouse cortical culture, CaEDTA selectively blocks  $Zn^{2+}$ -induced neuronal degeneration, but not the  $Ca^{2+}$ -overload neurotoxicity induced by glutamate agonists.

35 2) In a rat model of transient forebrain ischemia (bilateral carotid occlusion combined with hypotension), intraventricular injections of

CaEDTA, 30 minutes prior to the beginning of ischemia, reduced the ischemia-induced translocation of  $Zn^{2+}$  from presynaptic terminals throughout the brain to degenerating postsynaptic neurons. This CaEDTA treatment also markedly reduced the death of hippocampal hilar and CA1 pyramidal neurons.

### Example 1

#### Use of disodium-calcium-EDTA (CaEDTA) in a rat model of transient global ischemia.

5  $\mu$ l of 100 mM CaEDTA in saline, or saline alone, was injected into the lateral ventricles of Long Evans rats, 30 minutes prior to the inducement of 10 minutes of global ischemia. Ischemia was delivered by ligation of both common carotid arteries combined with hypotension after anesthesia (Smith et al., 1984).

Hippocampal sections from the treated and untreated rats were stained with the zinc-sensitive dye, TSQ (Frederickson et al., 1987). The examination of these sections showed that in the untreated rats zinc translocated from presynaptic inputs to degenerating hilar neuronal cell bodies between 3-24 hours after the ischemia. In the treated rats, CaEDTA blocked not only this zinc translocation, but also prevented subsequent hilar neuronal degeneration as assessed 3 days later.

TSQ staining also revealed the delayed appearance of zinc in CA1 pyramidal neuronal cell bodies of untreated rats 24-72 hours post ischemia. The CaEDTA treatment prior to the induced ischemia attenuated both the appearance of zinc in CA1 pyramidal neurons, and CA1 neuronal degeneration.

Control intraventricular injections of zinc EDTA, which is not a zinc chelating compound, did not alter the zinc translocation or hippocampal neuronal death following the ischemia in treated rats.

These results demonstrate that extracellular  $Zn^{2+}$  chelating compounds protect brain cells against neurotoxic injury which can be the result of ischemia or other neurodegenerative conditions associated with  $Zn^{2+}$  release.

Claims

1. The use of pharmaceutically acceptable zinc chelating compounds for the manufacture of medicaments for the treatment of neurotoxic injury.

2. The use of zinc chelating compounds according to claim 1, wherein  
5 the therapeutic indications include neurological insults such as ischemia as a result of stroke, cardiac arrest, hypoglycemia, epilepsy or trauma, neurodegenerative disorders such as Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and the neurodegenerative effects of AIDS.

10 3. The use of zinc chelating compounds according to claims 1 and 2, wherein the zinc chelating compounds are selected from the group consisting of  
disodium-calcium-ethylenediaminetetraacetic acid,  
3-mercapto-D-valine,  
15 (ethylenedioxy)diethylenedinitrilotetraacetic acid (EGTA),  
N,N,N',N'-tetrakis(2-pyridylmethyl)-ethylenediamine,  
N-(6-methoxy-8-quinolyl)-p-toluenesulfonamide,  
8-hydroxy quinoline,  
8-hydroxy quinoline-5-sulphonic acid,  
20 diethyl-dithiocarbamate,  
bis(diethylthiocarbamoyl)-disulfide,  
1,10-phenanthroline,  
dipicolinate,  
N-acetyl cystein,  
25 diphenylthiocarbazone,  
1-[2-(5-carboxyoxazol-2-yl)-6-aminobenzofuran-5-oxy]-2-(2'-amino-5'-methylphenoxy)ethane-N,N,N',N'-tetraacetic acid and  
1,2-bis(2-aminophenoxy)-ethane-N,N,N',N'-tetraacetic acid.

4. A medicament containing one or more zinc chelating compounds as  
30 defined in claim 3 and a pharmaceutically acceptable inert carrier for the treatment of neurotoxic injury which include neurological insults such as ischemia as a result of stroke, cardiac arrest, hypoglycemia, epilepsy or trauma, neurodegenerative disorders such as Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and the  
35 neurodegenerative effects of AIDS.




5. A method of treating neurotoxic injury in a patient suffering said injury by injecting or infusing into the cerebrospinal fluid of said patient a composition comprising a pharmaceutically acceptable zinc chelating compound as defined in claim 3 and a pharmaceutically acceptable carrier
- 5 wherein said zinc chelating compound is present in said composition in an amount sufficient to treat said neurotoxic injury.

6. The method according to claim 5, wherein the zinc chelating compound is administered at a dosage in the range from about 0,1 mg/kg to about 100 mg/kg daily.

10

\*\*\*





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>6</sup> :</b> <b>A61K 33/30, 31/13, 31/195, 31/325,</b> <b>31/44, 31/47, 31/42</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 97/09976</b> <b>(43) International Publication Date:</b> 20 March 1997 (20.03.97)
<b>(21) International Application Number:</b> PCT/IB96/00981 <b>(22) International Filing Date:</b> 23 August 1996 (23.08.96)  <b>(30) Priority Data:</b> 60/003,134 1 September 1995 (01.09.95) US 60/007,356 20 November 1995 (20.11.95) US  <b>(71) Applicant:</b> WASHINGTON UNIVERSITY [US/US]; One Brookings Drive, Campus Box 8013, St. Louis, MO 63130 (US). <b>(72) Inventors:</b> CHOI, Dennis, Wonkyu; 7400 Forsyth Boulevard, St. Louis, MO 63105 (US). KOH, Jae-young; 8148 Flintstone Trail, St. Louis, MO 63123 (US). <b>(74) Agent:</b> POKRAS, Bruce, A.; 340 Kingsland Street, Nutley, NJ 07110-1199 (US).	<b>(81) Designated States:</b> AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 22 May 1997 (22.05.97)	

**(54) Title:** METHOD OF REDUCING NEUROTOXIC INJURY WITH ZINC CHELATORS**(57) Abstract**

The invention relates to the use of pharmaceutically acceptable zinc chelating compounds for the manufacture of medicaments for the treatment of neurotoxic injury.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

# INTERNATIONAL SEARCH REPORT

Int. Application No.  
PCT/IB 96/00981

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K33/30 A61K31/13 A61K31/195 A61K31/325 A61K31/44  
A61K31/47 A61K31/42

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 93 10459 A (UNIV MELBOURNE) 27 May 1993 see page 6, line 13-20 see page 7, line 30 - page 8, line 2 see page 8, line 21 - page 9, line 14 ---	1-6
X	US 5 206 264 A (MARANGOS PAUL J) 27 April 1993 see the whole document ---	1-4
X	US 5 373 021 A (MARANGOS PAUL J) 13 December 1994 see the whole document ---	1-4
X	DE 39 32 338 A (NMI NATURWISSENSCHAFTL U MEDIZ) 11 April 1991 see column 2 ---	1,2
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*&\* document member of the same patent family

Date of the actual completion of the international search

7 April 1997

Date of mailing of the international search report

18.04.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Trifilieff-Riolo, S

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 96/00981

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 656 362 A (LILLY INDUSTRIES LTD) 7 June 1995 see page 9, line 4-10 see page 3, line 38-41 ---	1,2
X	DATABASE DISSERTATION ABSTRACTS umi aadaa-immo4720, 1995 LEE P.J.: "modulation of central cholinergic excitotoxicity by zinc chelating agents" XP002028935 see abstract	1-6
X	& MASTERS ABSTRACTS, vol. 34, no. 04, 1995, page 1564 LEE P.J.: "idem" see abstract ---	1-6
X,P	WO 96 06639 A (MEDICO PHARMA VERTRIEBS GMBH ; SICOMED S A (RO); BACANU SERBAN AL ( ) 7 March 1996 p.6 last paragraph; p.7 first paragraph; p.5 ex. 1-8 ---	1-4
X,P	SCIENCE, vol. 272, no. 5264, 17 May 1996, pages 1013-1016, XP002028933 KOH ET AL: "the role of zinc in selective neuronal death after transient global cerebral ischemia" see the whole document ---	1-6
X,P	P.N.A.S., vol. 93, no. 4, February 1996, pages 1710-1715, XP002028934 ARISPE ET AL: "zn2+ interaction with alzheimer amyloid beta protein calcium channels" see page 1713, right-hand column - page 1714, left-hand column ---	1-4
X,P	NEUROREPORT, vol. 7, no. 7, 17 May 1996, pages 1301-1304, XP000670010 CUAJUNGO ET AL: "prevention of zinc neurotoxicity in vivo by tpen" see the whole document ---	1-6

-/--

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 96/00981

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NEUROSCIENCE, vol. 60, no. 4, 1994, pages 1049-1057, XP000670025 KOH ET AL: "zinc toxicity on cultured cortical neurons" cited in the application see the whole document -----</p>	1-6

# INTERNATIONAL SEARCH REPORT

Information on patent family members

In .ional Application No

PCT/IB 96/00981

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9310459 A	27-05-93	AU 669493 B CA 2123211 A EP 0613560 A JP 7503316 T	13-06-96 27-05-93 07-09-94 06-04-95
US 5206264 A	27-04-93	CA 2094843 A US 5373021 A	27-10-94 13-12-94
US 5373021 A	13-12-94	CA 2094843 A US 5206264 A	27-10-94 27-04-93
DE 3932338 A	11-04-91	NONE	
EP 0656362 A	07-06-95	JP 7196664 A US 5580900 A US 5574158 A	01-08-95 03-12-96 12-11-96
WO 9606639 A	07-03-96	DE 4431175 A AU 3519495 A	11-04-96 22-03-96





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A61K 31/195, 31/47, 33/30, 31/325, 31/42, 31/13, 31/44</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/09976</b> <b>(43) International Publication Date:</b> 20 March 1997 (20.03.97)
<b>(21) International Application Number:</b> PCT/IB96/00981 <b>(22) International Filing Date:</b> 23 August 1996 (23.08.96)  <b>(30) Priority Data:</b> 60/003,134 1 September 1995 (01.09.95) US 60/007,356 20 November 1995 (20.11.95) US  <b>(71) Applicant:</b> WASHINGTON UNIVERSITY [US/US]; One Brookings Drive, Campus Box 8013, St. Louis, MO 63130 (US).  <b>(72) Inventors:</b> CHOI, Dennis, Wonkyu; 7400 Forsyth Boulevard, St. Louis, MO 63105 (US). KOH, Jae-young; 8148 Flintstone Trail, St. Louis, MO 63123 (US).  <b>(74) Agent:</b> POKRAS, Bruce, A.; 340 Kingsland Street, Nutley, NJ 07110-1199 (US).	<b>(81) Designated States:</b> AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title:</b> METHOD OF REDUCING NEUROTOXIC INJURY WITH ZINC CHELATORS  <b>(57) Abstract</b>  The invention relates to the use of pharmaceutically acceptable zinc chelating compounds for the manufacture of medicaments for the treatment of neurotoxic injury.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

Method of reducing neurotoxic injury with zinc chelators

The invention relates to the use of zinc chelating compounds in the treatment of neurotoxic injury.

It has been discovered that zinc chelating compounds protect  
5 hippocampal neurons from the neurotoxic effects of  $Zn^{2+}$  released during neurotoxic events, such as ischemia as a result of stroke or cardiac arrest. Thus, the present invention comprises a method of treating neurotoxic injury in a patient suffering said injury, comprising administering to said patient a zinc chelating compound in an amount sufficient to treat said  
10 neurotoxic injury.

The administration of the composition comprising the zinc chelating compound is carried out by injection or infusion of the composition into the patients cerebrospinal fluid.

Chelatable  $Zn^{2+}$  is present in large quantities in presynaptic vesicles of  
15 central excitatory neurons (Danscher et al., 1985; Frederickson et al., 1983), and released with synaptic activity or membrane depolarization (Assaf and Chung, 1984; Howell et al., 1984; Charton et al., 1985). Although the precise role of released synaptic zinc is not known, it blocks NMDA receptor-mediated current (Westbrook and Mayer, 1987; Peters et al., 1987; Christine  
20 and Choi, 1990) and GABA receptor-mediated current (Westbrook and Mayer, 1987), as well as voltage-dependent calcium channels (Winegar and Lansman, 1990). In addition, exposure to excessive extracellular  $Zn^{2+}$  is neurotoxic to cortical neurons (Choi et al., 1988), possibly mediated by  $Zn^{2+}$  influx in part mediated by ionotropic glutamate receptors and voltage-gated  
25 calcium channels (Weiss et al., 1993; Koh and Choi, 1994). Recently, it has been shown that zinc translocates into degenerating hippocampal hilar neurons after transient ischemia (Tonder et al., 1990) or kainate-induced seizures (Frederickson et al., 1989).

Treating neurotoxic injury within the meaning of the present invention  
30 means reducing the extent of damage to central neurons surrounding a central neuron which has released  $Zn^{2+}$  due to its having been damaged by a

neurotoxic event. Neurotoxic events include acute neurological insults such as hypoxia/ischemia, such as occurs during stroke, cardiac arrest, hypoglycemia, epilepsy or trauma. Neurotoxic events may also be chronic neuronal damage caused by neurodegenerative disorders such as

5 Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and the neurodegenerative effects of AIDS. Thus, the present invention also comprises a method of treating diseases, such as those described above, in which said neurotoxic injury occurs.

The zinc chelating compounds useful in accordance with the invention

10 are not critical. Any conventional compound which is capable of chelating  $Zn^{2+}$  and which is pharmaceutically acceptable for injection into cerebral spinal fluid may be used in accordance with the invention.

While ethylenediaminetetraacetic acid (EDTA) is traditionally used as an organic chelating agent for  $Ca^{2+}$ , it has affinity for other divalent metal

15 ions, and specifically binds  $Zn^{2+}$  with much higher affinity than  $Ca^{2+}$  (log stability constant at pH 7, 13.1 for  $Zn^{2+}$  versus 7.3 for  $Ca^{2+}$ ). An equimolar combination for  $Ca^{2+}$  and EDTA will act as a chelating agent for  $Zn^{2+}$  without chelating local  $Ca^{2+}$  (Dansher et al., 1975). Thus, the preferred zinc chelating compound for use in accordance with the invention is disodium-

20 calcium EDTA ("CaEDTA"), especially the form of CaEDTA known as edetate calcium disodium injection, USP (e.g., Calcium Disodium Versenate, 3M Pharmaceuticals, St. Paul, Minnesota).

Another preferred zinc chelating compound useful in accordance with the invention is 3-mercapto-D-valine (penicillamine), which is a chelating

25 agent usually applied to the treatment of Wilson's disease where it removes excess copper.

Other zinc chelating compounds useful in accordance with the invention are:

bis(diethylthiocarbamoyl) disulfide (Disulfiram or Antabuse),

30 (ethylenedioxy)diethylenedinitrilotetraacetic acid (EGTA),  
N,N,N',N'-tetrakis(2-pyridylmethyl)-ethylenediamine (TPEN),  
N-(6-methoxy-8-quinolyl)-p-toluenesulfonamide (TSQ),  
8-hydroxy quinoline (Oxyne),  
8-hydroxy quinoline-5-sulphonic acid (Sulphoxine),

35 diethyl dithiocarbamate (DEDTC),

- 1,10-phenanthroline,  
dipicolinate,  
N-acetyl cystein,  
diphenylthiocarbazone (Dithizone),  
5 1-[2-(5-carboxyoxazol-2-yl)-6-aminobenzofuran-5-oxy]-2-(2'-amino-5'-methylphenoxy)ethane-N,N,N',N'-tetraacetic acid (Fura-2), and  
1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (BAPTA).

The zinc chelating compound used in accordance with the present invention may be administered to the cerebrospinal fluid by any conventional  
10 means. The preferred method of administration is by injection or chronic pump infusion into the lateral ventricles of the patient's brain or into the lumbar sac of the patient.

The zinc chelating compound is preferably administered as a composition containing the zinc chelating compound and a  
15 pharmaceutically acceptable carrier compatible with the compound. In preparing such a composition, any conventional pharmaceutically acceptable carrier may be utilized. Typical carriers for administration by injection would be sterile aqueous/buffered solutions, preferably water for injection or unbuffered or buffered physiological saline. The zinc chelating  
20 compound is preferably present in the carrier at a concentration of 1 mM - 300 mM, especially from 50 mM - 200 mM.

In carrying out the method of the invention, the zinc chelating compound is administered to adults daily in an amount from about 0.1 mg/kg to about 100 mg/kg daily, in single or divided doses, or continuously  
25 through chronic pump infusion. The preferred dosage will vary depending upon the indication for which the method of the invention is being used to treat. For treatment of patients who have suffered an acute neurotoxic insult, as a result of, e.g., stroke or cardiac arrest, the administration of a dosage from about 5 mg/kg to about 50 mg/kg daily, carried out for from 1 to 7  
30 days, is preferred. Administration may be carried out by injection or by infusion utilizing a chronic infusion pump. For treatment of chronic neurotoxic conditions in patients, e.g., Alzheimer's disease, a dosage of from about 1 mg/kg to about 10 mg/kg daily is preferred, carried out for months to years, preferably utilizing a chronic infusion pump.

35 Thus, the present invention also comprises a method of treating neurotoxic injury in a patient suffering said injury by administering to said

patient a composition comprising a zinc chelating compound and carrier, wherein both the compound and the carrier are pharmaceutically acceptable for injection. The preferred zinc chelating compounds are CaEDTA and 3-mercapto-D-valine, with CaEDTA being especially preferred. The zinc  
5 chelating compound is preferably administered in an amount from about 0.1 mg/kg to about 100 mg/kg daily, especially in an amount from about 5 mg/kg to about 50 mg/kg daily for the treatment of acute neurotoxic conditions and in an amount from about 1 mg/kg to about 10 mg/kg daily for chronic  
10 neurotoxic conditions. The preferred method of administration is the injection or infusion of the pharmaceutical composition comprising the zinc chelating compound and the carrier into the cerebrospinal fluid (e.g., the lateral ventricles or lumbar sac) of the patient. The preferred duration of treatment is from 1 to 7 days for acute neurotoxic conditions, and for months to years for chronic neurotoxic conditions.

15 The present invention preferably comprises a method of treating stroke in a patient suffering said stroke by administering to said patient a composition comprising a zinc chelating compound and a carrier, wherein both the compound and the carrier are pharmaceutically acceptable for injection. The preferred zinc chelating compounds are CaEDTA and 3-  
20 mercapto-D-valine, with CaEDTA being especially preferred. The zinc chelating compound is preferably administered in an amount from about 0.1 mg/kg to about 100 mg/kg daily, especially in an amount from about 5 mg/kg to about 50 mg/kg daily for the treatment of stroke. The preferred method of administration is the injection or infusion, especially injection, of the  
25 pharmaceutical composition comprising the zinc chelating compound and the carrier into the cerebrospinal fluid (e.g., the lateral ventricles or lumbar sac) of the patient. The preferred duration of treatment is from 1 to 7 days.

Experiments were conducted which determined that a zinc chelating compound, CaEDTA, exhibited neuroprotective effects in cell culture and  
30 animal models of brain hypoxic-ischemic injury, such as might occur consequent to cardiac arrest or stroke. Several observations were made:

1) In mouse cortical culture, CaEDTA selectively blocks  $Zn^{2+}$ -induced neuronal degeneration, but not the  $Ca^{2+}$ -overload neurotoxicity induced by glutamate agonists.

35 2) In a rat model of transient forebrain ischemia (bilateral carotid occlusion combined with hypotension), intraventricular injections of

CaEDTA, 30 minutes prior to the beginning of ischemia, reduced the ischemia-induced translocation of  $Zn^{2+}$  from presynaptic terminals throughout the brain to degenerating postsynaptic neurons. This CaEDTA treatment also markedly reduced the death of hippocampal hilar and CA1  
5 pyramidal neurons.

### Example 1

#### Use of disodium-calcium-EDTA (CaEDTA) in a rat model of transient global ischemia.

5  $\mu$ l of 100 mM CaEDTA in saline, or saline alone, was injected into the  
10 lateral ventricles of Long Evans rats, 30 minutes prior to the inducement of 10 minutes of global ischemia. Ischemia was delivered by ligation of both common carotid arteries combined with hypotension after anesthesia (Smith et al., 1984).

Hippocampal sections from the treated and untreated rats were stained  
15 with the zinc-sensitive dye, TSQ (Frederickson et al., 1987). The examination of these sections showed that in the untreated rats zinc translocated from presynaptic inputs to degenerating hilar neuronal cell bodies between 3-24 hours after the ischemia. In the treated rats, CaEDTA blocked not only this zinc translocation, but also prevented subsequent hilar neuronal  
20 degeneration as assessed 3 days later.

TSQ staining also revealed the delayed appearance of zinc in CA1 pyramidal neuronal cell bodies of untreated rats 24-72 hours post ischemia. The CaEDTA treatment prior to the induced ischemia attenuated both the appearance of zinc in CA1 pyramidal neurons, and CA1 neuronal  
25 degeneration.

Control intraventricular injections of zinc EDTA, which is not a zinc chelating compound, did not alter the zinc translocation or hippocampal neuronal death following the ischemia in treated rats.

These results demonstrate that extracellular  $Zn^{2+}$  chelating  
30 compounds protect brain cells against neurotoxic injury which can be the result of ischemia or other neurodegenerative conditions associated with  $Zn^{2+}$  release.

Claims

1. The use of pharmaceutically acceptable zinc chelating compounds for the manufacture of medicaments for the treatment of neurotoxic injury.

2. The use of zinc chelating compounds according to claim 1, wherein  
5 the therapeutic indications include neurological insults such as ischemia as a result of stroke, cardiac arrest, hypoglycemia, epilepsy or trauma, neurodegenerative disorders such as Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and the neurodegenerative effects of AIDS.

10 3. The use of zinc chelating compounds according to claims 1 and 2, wherein the zinc chelating compounds are selected from the group consisting of

disodium-calcium-ethylenediaminetetraacetic acid,  
3-mercapto-D-valine,  
15 (ethylenedioxy)diethylenedinitrilotetraacetic acid (EGTA),  
N,N,N',N'-tetrakis(2-pyridylmethyl)-ethylenediamine,  
N-(6-methoxy-8-quinolyl)-p-toluenesulfonamide,  
8-hydroxy quinoline,  
8-hydroxy quinoline-5-sulphonic acid,  
20 diethyl-dithiocarbamate,  
bis(diethylthiocarbamoyl)-disulfide,  
1,10-phenanthroline,  
dipicolinate,  
N-acetyl cystein,  
25 diphenylthiocarbazone,  
1-[2-(5-carboxyoxazol-2-yl)-6-aminobenzofuran-5-oxy]-2-(2'-amino-5'-methylphenoxy)ethane-N,N,N',N'-tetraacetic acid and  
1,2-bis(2-aminophenoxy)-ethane-N,N,N',N'-tetraacetic acid.

4. A medicament containing one or more zinc chelating compounds as  
30 defined in claim 3 and a pharmaceutically acceptable inert carrier for the treatment of neurotoxic injury which include neurological insults such as ischemia as a result of stroke, cardiac arrest, hypoglycemia, epilepsy or trauma, neurodegenerative disorders such as Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis, and the  
35 neurodegenerative effects of AIDS.



5. A method of treating neurotoxic injury in a patient suffering said injury by injecting or infusing into the cerebrospinal fluid of said patient a composition comprising a pharmaceutically acceptable zinc chelating compound as defined in claim 3 and a pharmaceutically acceptable carrier
- 5 wherein said zinc chelating compound is present in said composition in an amount sufficient to treat said neurotoxic injury.

6. The method according to claim 5, wherein the zinc chelating compound is administered at a dosage in the range from about 0,1 mg/kg to about 100 mg/kg daily.





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A61K 33/30, 31/13, 31/195, 31/325,</b> <b>31/44, 31/47, 31/42</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 97/09976</b> <b>(43) International Publication Date:</b> 20 March 1997 (20.03.97)
<b>(21) International Application Number:</b> PCT/IB96/00981 <b>(22) International Filing Date:</b> 23 August 1996 (23.08.96)  <b>(30) Priority Data:</b> 60:003,134 1 September 1995 (01.09.95) US 60:007,356 20 November 1995 (20.11.95) US  <b>(71) Applicant:</b> WASHINGTON UNIVERSITY [US/US]; One Brookings Drive, Campus Box 8013, St. Louis, MO 63130 (US). <b>(72) Inventors:</b> CHOI, Dennis, Wonkyu; 7400 Forsyth Boulevard, St. Louis, MO 63105 (US). KOH, Jae-young; 8148 Flintstone Trail, St. Louis, MO 63123 (US). <b>(74) Agent:</b> POKRAS, Bruce, A.; 340 Kingsland Street, Nutley, NJ 07110-1199 (US).	<b>(81) Designated States:</b> AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 22 May 1997 (22.05.97)	
<b>(54) Title:</b> METHOD OF REDUCING NEUROTOXIC INJURY WITH ZINC CHELATORS  <b>(57) Abstract</b>  The invention relates to the use of pharmaceutically acceptable zinc chelating compounds for the manufacture of medicaments for the treatment of neurotoxic injury.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## INTERNATIONAL SEARCH REPORT

Int. l. Application No.

PCT/IB 96/00981

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K33/30 A61K31/13 A61K31/195 A61K31/325 A61K31/44  
 A61K31/47 A61K31/42

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 93 10459 A (UNIV MELBOURNE) 27 May 1993 see page 6, line 13-20 see page 7, line 30 - page 8, line 2 see page 8, line 21 - page 9, line 14 ---	1-6
X	US 5 206 264 A (MARANGOS PAUL J) 27 April 1993 see the whole document ---	1-4
X	US 5 373 021 A (MARANGOS PAUL J) 13 December 1994 see the whole document ---	1-4
X	DE 39 32 338 A (NMI NATURWISSENSCHAFTL U MEDIZ) 11 April 1991 see column 2 --- -/-	1,2

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*A\* document member of the same patent family

Date of the actual completion of the international search

7 April 1997

Date of mailing of the international search report

18.04.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+31-70) 340-3016

Authorized officer

Trifilieff-Riolo, S

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 96/00981

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 656 362 A (LILLY INDUSTRIES LTD) 7 June 1995 see page 9, line 4-10 see page 3, line 38-41 ---	1,2
X	DATABASE DISSERTATION ABSTRACTS umi aadaa-immo4720, 1995 LEE P.J.: "modulation of central cholinergic excitotoxicity by zinc chelating agents" XP002028935 see abstract	1-6
X	& MASTERS ABSTRACTS, vol. 34, no. 04, 1995, page 1564 LEE P.J.: "idem" see abstract ---	1-6
X,P	WO 96 06639 A (MEDICO PHARMA VERTRIEBS GMBH ; SICOMED S A (RO); BACANU SERBAN AL ( ) 7 March 1996 p.6 last paragraph; p.7 first paragraph; p.5 ex. 1-8 ---	1-4
X,P	SCIENCE, vol. 272, no. 5264, 17 May 1996, pages 1013-1016, XP002028933 KOH ET AL: "the role of zinc in selective neuronal death after transient global cerebral ischemia" see the whole document ---	1-6
X,P	P.N.A.S., vol. 93, no. 4, February 1996, pages 1710-1715, XP002028934 ARISPE ET AL: "zn2+ interaction with alzheimer amyloid beta protein calcium channels" see page 1713, right-hand column - page 1714, left-hand column ---	1-4
X,P	NEUROREPORT, vol. 7, no. 7, 17 May 1996, pages 1301-1304, XP000670010 CUAJUNGO ET AL: "prevention of zinc neurotoxicity in vivo by tpen" see the whole document ---	1-6
	-/--	

2

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 96/00981

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NEUROSCIENCE,  vol. 60, no. 4, 1994,  pages 1049-1057, XP000670025  KOH ET AL: "zinc toxicity on cultured  cortical neurons"  cited in the application  see the whole document  -----</p>	1-6

Form PCT-ISA/210 (continuation of second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/IB 96/00981

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9310459 A	27-05-93	AU 669493 B CA 2123211 A EP 0613560 A JP 7503316 T	13-06-96 27-05-93 07-09-94 06-04-95
US 5206264 A	27-04-93	CA 2094843 A US 5373021 A	27-10-94 13-12-94
US 5373021 A	13-12-94	CA 2094843 A US 5206264 A	27-10-94 27-04-93
DE 3932338 A	11-04-91	NONE	
EP 0656362 A	07-06-95	JP 7196664 A US 5580900 A US 5574158 A	01-08-95 03-12-96 12-11-96
WO 9606639 A	07-03-96	DE 4431175 A AU 3519495 A	11-04-96 22-03-96